

Problem 1 (Legendre transform, 2+2+2 points).

For a function $H : \mathbb{R}^n \rightarrow \mathbb{R}$, denote by $L = H^*$ its Legendre transform.

- a) Let $H(p) := \frac{1}{r}|p|^r$, for $1 < r < \infty$. Show that $L(q) = \frac{1}{r'}|q|^{r'}$, where $\frac{1}{r} + \frac{1}{r'} = 1$.
- b) Let $H(p) := \frac{1}{2}p \cdot Ap + b \cdot p$, where $A \in \mathbb{R}^{n \times n}$ is a symmetric, positive definite matrix, and $b \in \mathbb{R}^n$. Compute L .
- c) Let $H(p) = \sqrt{1 + p^2}$, $p \in \mathbb{R}$. Compute L .

Problem 2 (Subdifferential and Hopf-Lax formula, 2+2 points).

Let $H : \mathbb{R}^n \rightarrow \mathbb{R}$ be a convex function.

- a) We say that $q \in \mathbb{R}^n$ belongs to the subdifferential of H at p , written $q \in \partial H(p)$, if

$$H(r) \geq H(p) + q \cdot (r - p) \quad \text{for all } r \in \mathbb{R}^n.$$

Prove that

$$q \in \partial H(p) \iff p \in \partial L(q) \iff p \cdot q = H(p) + L(q),$$

where $L = H^*$ is the Legendre transform of H .

- b) Assume also that H and g are of class C^1 and $\lim_{|p| \rightarrow \infty} \frac{H(p)}{|p|} = \infty$. Prove that the Hopf-Lax formula reads

$$u(x, t) = \min_{y \in \mathbb{R}^n} \left\{ tL\left(\frac{x-y}{t}\right) + g(y) \right\} = \min_{y \in B(x, Rt)} \left\{ tL\left(\frac{x-y}{t}\right) + g(y) \right\}$$

for $R = \sup_{y \in \mathbb{R}^n} |DH(Dg(y))|$, $H = L^*$. Here $B(x, Rt) = \{y \in \mathbb{R}^n : |y - x| < Rt\}$. (This proves *finite propagation speed* for Hamilton-Jacobi equation).

Hint: you can use the fact that, for a convex, C^1 function H , the subdifferential $\partial H(p)$ contains precisely one element, $DH(p)$.

Problem 3 (Hopf-Lax formula, 2 points).

Use Hopf-Lax formula to solve explicitly the PDE

$$\begin{cases} u_t + \frac{3}{4}u^{\frac{4}{3}} = 0 & \text{for } (x, t) \in \mathbb{R} \times (0, \infty), \\ u(x, 0) = x. \end{cases}$$

Problem 4 (L^∞ -contraction property of Hamilton-Jacobi equations, 4 points).

Let $H \in C^2(\mathbb{R}^n)$ be uniformly convex with $\lim_{|p| \rightarrow \infty} \frac{H(p)}{|p|} = \infty$, and let $g_1, g_2 : \mathbb{R}^n \rightarrow \mathbb{R}$ be Lipschitz continuous. Assume u^1, u^2 are weak solutions of the initial-value problem

$$\begin{cases} u_t^i + H(Du^i) = 0 & \text{in } \mathbb{R}^n \times (0, \infty), \\ u^i = g^i & \text{on } \mathbb{R}^n \times \{t = 0\}, \end{cases}$$

$i = 1, 2$. Prove the inequality

$$\|u^1(\cdot, t) - u^2(\cdot, t)\|_{L^\infty(\mathbb{R}^n)} \leq \|g^1 - g^2\|_{L^\infty(\mathbb{R}^n)} \quad \text{for all } t > 0.$$

Total: 16 points