

Problem 1 (Macroscopic balance law).

Suppose that $f = f(t, x, \xi)$ is a solution to Boltzmann equation. Define mass and momentum density as follows:

$$\rho(t, x) = \int_{\mathbb{R}^3} f(t, x, \xi) \, d\xi,$$

$$(\rho u)(t, x) = \int_{\mathbb{R}^3} \xi f(t, x, \xi) \, d\xi.$$

Furthermore let $c = \xi - u \in \mathbb{R}^3$ and define the internal energy density and the heat and momentum flow as

$$(\rho e)(t, x) = \frac{1}{2} \int_{\mathbb{R}^3} |c|^2 f(t, x, \xi) \, d\xi,$$

$$q_i(t, x) = \frac{1}{2} \int_{\mathbb{R}^3} c_i |c|^2 f(t, x, \xi) \, d\xi,$$

$$p_{ij}(t, x) = \int_{\mathbb{R}^3} c_i c_j f(t, x, \xi) \, d\xi,$$

$i, j = 1, 2, 3$. Show that the following equations hold:

$$\begin{cases} \partial_t \rho + \operatorname{div}(\rho u) = 0, \\ \partial_t (\rho u_j) + \operatorname{div}(\rho u_j u + p_{.j}) = 0, \\ \partial_t \left(\frac{1}{2} \rho |u|^2 + \rho e \right) + \operatorname{div} \left(\rho u \left(\frac{1}{2} |u|^2 + e \right) \right) + \operatorname{div}(p u + q) = 0. \end{cases} \quad j = 1, 2, 3,$$

What do you obtain if f is independent of x ?

Hint: choose suitable collision invariants.

Problem 2 (Maxwell distributions).

Let $f = f(t, \xi)$ be a spatially homogeneous solution to Boltzmann equation (that is, f does not depend on the spatial variable x). Recall that in this case the quantities

$$\rho(t) := \int_{\mathbb{R}^3} f(t, \xi) \, d\xi, \quad (\rho u)(t) := \int_{\mathbb{R}^3} \xi f(t, \xi) \, d\xi, \quad (\rho e)(t) := \frac{1}{2} \int_{\mathbb{R}^3} |\xi|^2 f(t, \xi) \, d\xi$$

are constant in time. Show that, if M is the Maxwellian with the same $\rho, \rho u, \rho e$ as f , then

$$\mathcal{H}(f) \geq \mathcal{H}(M), \quad \text{where } \mathcal{H}(f) := \int_{\mathbb{R}^3} f \ln f \, d\xi.$$

Hint: first show that $\int_{\mathbb{R}^3} \ln M (f - M) \, d\xi = 0$, using the fact that $\ln M$ is a collision invariant; moreover, use the inequality $z \ln z - z \ln y + y - z \geq 0$.