

YWIP 2014 - 27th May 2014
Mathematik Zentrum, Bonn

Static fluctuations of a thick $1D$ interface in the $1+1$ Directed Polymer formulation

Elisabeth Agoritsas

(LIPhy, Université Joseph Fourier de Grenoble)

Collaboration with Vivien Lecomte (LPMA - Paris VI-VII)
& Prof. Thierry Giamarchi (Université de Genève)



UNIVERSITÉ
DE GENÈVE

FACULTÉ DES SCIENCES
Département de physique
de la matière condensée



FONDS NATIONAL SUISSE
SCHWEIZERISCHER NATIONALFONDS
FONDO NAZIONALE SVIZZERO
SWISS NATIONAL SCIENCE FOUNDATION




MaNEP
SWITZERLAND



Laboratoire
interdisciplinaire de Physique

Disordered medium



Interface →

Interface width ↔

Interfaces can be found everywhere...



← Scotland coastline


Lengthscale matters!



← Crack in a pavement


Scale invariance?

Crack at the Moon's surface →




Interfaces can be found everywhere...


Magnetic thin film




Lemerle et al., *Phys. Rev. Lett.* **80**, 894 (1998).




Moulinet et al., *Eur. Phys. J. E* **8**, 437 (2002).



Crack fronts



Buldyrev et al., *Phys. Rev. A* **45**, 8313 (1992).



← Paper

← Sandblasted plexiglass

Santucci et al., *Phys. Rev. E* **75**, 016104 (2007).


Interfaces can be found everywhere...



Wetting line




Burning front




Crack fronts

Magnetic thin film

Lemerle et al., *Phys. Rev. Lett.* **80**, 894 (1998).




Imbibition front



(a)

← Paper



(b)


← Sandblasted plexiglass

Santucci et al., *Phys. Rev. E* **75**, 016104 (2007).

Interfaces can be found everywhere...


- Ubiquitous in Nature, large variety of lengthscales & microphysics.

BUT do they share nevertheless common (universal?) features?




Review: A.-L. Barabási & H. E. Stanley, *Fractal Concepts in Surface Growth*, Cambridge University Press, 1995.

- Increasing complexity starting from a **MICROSCOPIC** description.
⇒ Need of a simpler **MESOSCOPIC** starting point
- Systems supported by an inhomogeneous underlying medium.
⇒ Statistical characterization of **DISORDER**
- Effective description depending on the **LENGTHSCALE**.
⇒ Characteristic lengthscales, scale invariance?



How do they look like?
How do they respond when one pulls at them?
Disorder-conditioned features?





Disordered Elastic Systems (DES)

- Competition of three physical ingredients \Rightarrow METASTABILITY, GLASSY PROPERTIES

ELASTICITY

DISORDER

TEMPERATURE




Disordered Elastic Systems (DES)

- Competition of three physical ingredients \Rightarrow METASTABILITY, GLASSY PROPERTIES



ELASTICITY

DISORDER

TEMPERATURE



- Exploration of disordered energy landscapes



Disordered Elastic Systems (DES): a recipe

■ Dimensionality

■ **Elasticity:** Short-range versus long-range, e.g.



$$\mathcal{H}_{el} \propto \text{system size}$$

■ **Disorder:** - Quenched versus annealed disorder

- 'Random-bond' versus 'random-field'


- Collective weak pinning versus strong individual pinning centers

$$\mathcal{H}_{DES} = \mathcal{H}_{el} + \mathcal{H}_{dis}$$




■ **No bubbles nor overhangs**


■ **Finite width / Disorder correlation**



■ **Internal degree of freedom?**



What is the imprint of a finite microscopic width and/or disorder correlation length ξ on the 1D interface fluctuations and properties?




Main issue: finite width or disorder correlation length $\xi > 0$

- Two examples of experimental realizations of interfaces:

Ferromagnetic domain wall ($\xi \sim 50\text{nm}$)

RESOLUTION: $1\mu\text{m}$

Ultrathin film of Pt/Co/Pt (a few atomic layers)




S. Lemerle, J. Ferré, C. Chappert, V. Mathet, T. Giamarchi, & P. Le Doussal,
Phys. Rev. Lett. **80**, 849 (1998).

Ferroelectric domain wall ($\xi \sim 1\text{nm}$)

RESOLUTION: 5nm



PbZr_{0.2}Ti_{0.8}O₃ 70nm / SrRuO₃ 30nm (electrode) /
SrTiO₃ (substrate)



Courtesy of J. Guyonnet & Prof. P. Paruch.

Main result: low-temperature regime at $\xi > 0$

Random potential: $V(z, x)$





Interplay between
Thermal fluctuations $T > 0$
Width and/or disorder
correlation length $\xi > 0$



$$\xi_{\text{thermal}}(T) \lesssim \xi$$



$$T \lesssim T_c(\xi)$$



Outline

■ Introduction

- Generic framework: Disordered Elastic Systems (DES)
- Specific issue: role a finite width or disorder correlation length

■ Model


- Geometrical fluctuations and roughness
- DES model of a one-dimensional (1D) interface
- Static 1D interface versus 1+1 Directed Polymer (DP)

■ Our results: Temperature-dependent fluctuations

- Disorder free-energy fluctuations
- Roughness: temperature-induced crossover


■ Conclusion & Perspectives

Geometrical fluctuations & roughness



- Lengthscale r
- Relative displacement $\Delta u(r)$

- Probability distribution function $\mathcal{P}(\Delta u(r))$




Roughness exponent ζ
 =
Signature of the predominant physics

- Roughness function

$$B(r) = \overline{\langle \Delta u(r)^2 \rangle} \sim A \cdot r^{2\zeta}$$


- Roughness exponent:

$$\begin{cases} \zeta_{\text{thermal}} = 1/2 \\ \zeta_{\text{KPZ}} = 2/3 \end{cases}$$




Geometrical fluctuations & roughness: experimental examples

Domain walls in ultrathin Pt/Co/Pt ferromagnetic films




$$\zeta_{\text{exp}} = 0.69 \pm 0.07$$




S. Lemerle et al., *Phys. Rev. Lett.* **80**, 849 (1998).

Fluid invasion in a porous medium



$$\zeta_{\text{exp}} = 0.63 \pm 0.04$$



Buldyrev et al., *Phys. Rev. A* **45**, 8313 (1992).



Issues regarding the roughness at $\xi > 0$

- How many roughness regimes ?
Characteristic crossover lengthscales ?

- Universal roughness amplitude ?

$$B(r, c, D, T, \xi) \sim A_{(c, D, T, \xi)} \cdot r^{2\zeta}$$

- Imprint of the disorder correlator $R_\xi(x)$?



Model of a **thick** 1D interface & 1+1 Directed Polymer (DP)

- Short-range elasticity & Elastic limit / Quenched random-bond weak disorder

Hamiltonian: $\mathcal{H} [u, \tilde{V}] = \int_{\mathbb{R}} dz \cdot \left[\frac{c}{2} (\nabla_z u_z)^2 + \underbrace{\int_{\mathbb{R}} dx \cdot \rho_{\xi}(x - u_z) \tilde{V}(z, x)}_{V(z, u_z)} \right]$


- Density $\rho_{\xi}(x - u_z)$ & random potential $\tilde{V}(z, u_z)$

$$\overline{\tilde{V}(z, x)} = 0$$

$$\overline{\tilde{V}(z, x) \tilde{V}(z', x')} = D \cdot \delta_{(z-z')} \delta_{(x-x')}$$

- Alternative: *correlated* effective potential $V(z, u_z)$


$$\overline{V(z, x) V(z', x')} = D \cdot \delta_{(z-z')} R_{\xi}(x - x')$$




Elastic constant c / Width ξ / Disorder strength D / Temperature T

Static 1D interface & Growing 1+1 Directed Polymer (DP)

- **Observable:** static geometrical fluctuations $\mathcal{P}(\Delta u(r))$ & roughness $B(r) = \overline{\langle \Delta u(r)^2 \rangle}$
 & Effective disorder experienced by the 1D interface at a given lengthscale r
 \leftrightarrow at fixed growing DP 'time' t



Integrating the thermal fluctuations
at short-'times'/lengthscales!





**1D interface
Lengthscale**

**Directed Polymer
Growing 'time'**

Static 1D interface & Growing 1+1 Directed Polymer (DP)

- **Observable:** static geometrical fluctuations $\mathcal{P}(\Delta u(r))$ & roughness $B(r) = \overline{\langle \Delta u(r)^2 \rangle}$
- & Effective disorder experienced by the 1D interface at a given lengthscale r
- \leftrightarrow at fixed growing DP 'time' t



$$\mathcal{P}_V(t, y) \propto e^{-[F_{\text{th}}(t, y) + \bar{F}_V(t, y)]/T}$$


Integrating the thermal fluctuations at short-'times'/lengthscales!



'Time-dependent free-energy landscape

Static 1D interface & Growing 1+1 Directed Polymer (DP)

- KPZ evolution equation for the total free-energy with 'sharp wedge' initial condition:



D. Huse, C. L. Henley & D. S. Fisher, *Phys. Rev. Lett.* **55** 2924 (1985).

M. Kardar, G. Parisi & Y.-C. Zhang, *Phys. Rev. Lett.* **56** 889 (1986).


$$\begin{cases} \partial_t F_V(t, y) = \frac{T}{2c} \partial_y^2 F_V(t, y) - \frac{1}{2c} [\partial_y F_V(t, y)]^2 + V(t, y) \\ \mathcal{P}_V(0, y) = e^{-F_V(0, y)/T} = \delta(y) \end{cases}$$

- Tilted KPZ equation for the disorder contribution to the free-energy:

E. Agoritsas, V. Lecomte & T. Giamarchi, *Phys. Rev. E* **87**, 042406 & 062405 (2013).

$$\begin{cases} \partial_t \bar{F}_V(t, y) = \frac{T}{2c} \partial_y^2 \bar{F}_V(t, y) - \frac{1}{2c} [\partial_y \bar{F}_V(t, y)]^2 - \frac{y}{t} \partial_y \bar{F}_V(t, y) + V(t, y) \\ \bar{F}_V(0, y) \equiv 0 \quad (\text{'flat' initial condition}) \end{cases}$$

Static 1D interface & Growing 1+1 Directed Polymer (DP)



Focus on the unknown part of the free-energy

Translation-invariant distribution:

$$\bar{\mathcal{P}} [\bar{F}_V(t, y + Y)] = \bar{\mathcal{P}} [\bar{F}_V(t, y)]$$

Starting point of numerical/analytical study

- Tilted KPZ equation for the disorder contribution to the free-energy:

*E. Agoritsas, V. Lecomte & T. Giamarchi, Phys. Rev. E **87**, 042406 & 062405 (2013).*


$$\begin{cases} \partial_t \bar{F}_V(t, y) = \frac{T}{2c} \partial_y^2 \bar{F}_V(t, y) - \frac{1}{2c} [\partial_y \bar{F}_V(t, y)]^2 - \frac{y}{t} \partial_y \bar{F}_V(t, y) + V(t, y) \\ \bar{F}_V(0, y) \equiv 0 \quad (\text{'flat' initial condition}) \end{cases}$$

Kardar-Parisi-Zhang (KPZ) equation

M. Kardar, G. Parisi & Y.-C. Zhang, « Dynamical Scaling of Growing Interfaces », *Phys. Rev. Lett.* **56** 889 (1986).

- Model for the time-evolution of the profile of a growing interface $h(t, \vec{x}) \leftrightarrow F_V(t, y)$

$$\partial_t h(t, \vec{x}) = \underbrace{\nu \nabla_{\vec{x}}^2 h(t, \vec{x})}_{\text{relaxation or diffusion}} + \underbrace{\frac{\lambda}{2} [\nabla_{\vec{x}} h(t, \vec{x})]^2}_{\text{slope-dependent lateral growth}} + \underbrace{\eta(t, \vec{x})}_{\text{random noise}}$$



- Gaussian statistical distribution of a **white** noise...

$$\begin{cases} \overline{\eta(t, \vec{x})} = 0 \\ \overline{\eta(t, \vec{x})\eta(t', \vec{x}')} = D \cdot \delta(t - t') \cdot \delta^{(d)}(\vec{x} - \vec{x}') \end{cases}$$

... and of a **colored** noise in 1D (d=1)

$$\overline{\eta(t, x)\eta(t', x')} = D \cdot \delta(t - t') \cdot R_{\xi}(x - x')$$

Kardar-Parisi-Zhang (KPZ) equation

M. Kardar, G. Parisi & Y.-C. Zhang, « Dynamical Scaling of Growing Interfaces », *Phys. Rev. Lett.* **56** 889 (1986).

- Model for the time-evolution of the profile of a growing interface $h(t, \vec{x}) \leftrightarrow F_V(t, y)$

$$\partial_t h(t, \vec{x}) = \underbrace{\nu \nabla_{\vec{x}}^2 h(t, \vec{x})}_{\text{relaxation or diffusion}} + \underbrace{\frac{\lambda}{2} [\nabla_{\vec{x}} h(t, \vec{x})]^2}_{\text{slope-dependent lateral growth}} + \underbrace{\eta(t, \vec{x})}_{\text{random noise}}$$

relaxation or
diffusion

slope-dependent lateral
growth

random
noise

Initial condition
 $h(0, \vec{x})$



Distribution of the
random noise
 $\bar{\mathcal{P}}[\eta(t, \vec{x})]$

- 1D KPZ universality class encompasses a wide range of problems:

Random matrices, Burgers equation in hydrodynamics, roughening phenomena & stochastic growth,

1+1 Directed Polymer (DP), our one-dimensional interface, ...

Fluctuations with power-law of exponent $\zeta_{\text{KPZ}} = 2/3$


J. Quastel, « Introduction to KPZ », CMD 2011, <http://www.math.toronto.edu/quastel/survey.pdf>.

Ivan Corwin, « The Kardar-Parisi-Zhang equation and universality class », <http://arxiv.org/abs/1106.1596>.

Issues regarding the roughness at $\xi > 0$

1D interface - Lengthscale


- How many roughness regimes?
Characteristic crossover lengthscales?
- Universal roughness amplitude?
 $B(t, c, D, T, \xi) \sim A_{(c, D, T, \xi)} \cdot t^{2\zeta}$
- Imprint of the disorder correlator $R_\xi(y)$?



$$F_{\text{th}}(t, y) + \bar{F}_V(t, y)$$

$$\approx F_{\text{th}}(t, y)$$

$$\approx \bar{F}_V(t, y)$$



1+1 Directed Polymer - 'Time'

- Fluctuations of disorder free-energy $\bar{F}_V(t, y)$
- Focus on its two-point correlator $\bar{R}(t, y)$
- Scaling of free-energy correlator amplitude
 $\tilde{D}_\infty(T, \xi)$

■ Free-energy two-point correlators:

$$\begin{cases} \bar{C}(t, y) \equiv \overline{[\bar{F}_V(t, y) - \bar{F}_V(t, 0)]^2} \\ \bar{R}(t, y) \equiv \overline{\partial_y \bar{F}_V(t, y) \partial_y \bar{F}_V(t, 0)} \end{cases}$$

■ Uncorrelated disorder (white-noise): $R_{\xi=0}(y) = \delta(y)$

■ Infinite-‘time’ limit:

$$\begin{cases} \text{Gaussian distribution} \\ \bar{C}(\infty, y) = \frac{cD}{T} |y| \iff \bar{R}(\infty, y) = \frac{cD}{T} R_{\xi=0}(y) \end{cases}$$

D.A. Huse, C. L. Henley & D. S. Fisher, *Phys. Rev. Lett.* **55** 2294 (1985).

■ Asymptotically large-‘time’:

$$\begin{cases} \text{GUE Tracy-Widom distribution (non-Gaussian!)} \\ \bar{C}(t, y) = \text{2-point correlator of Airy}_2 \text{ process} \end{cases}$$

M. Prähofer & H. Spohn, *J. Stat. Phys.* **159** 1071 (2002).

- At all ‘times’:
- P. Calabrese, P. Le Doussal & A. Rosso, *Eur. Phys. Lett.* **90** 20002 (2010).
 - V. Dotsenko, *Eur. Phys. Lett.* **90** 20003 (2010).
 - T. Sasamoto & H. Spohn, *Nucl. Phys. B* **834** 523 (2010).
 - G. Amir, I. Corwin, J. Quastel., *Comm. Pure Appl. Math.* **64** 466 (2011).

Free-energy of the $1+1$ DP: 'time'-dependence

$(\xi > 0)$



Free-energy two-point correlators:

$$\begin{cases} \bar{C}(t, y) \equiv \overline{[\bar{F}_V(t, y) - \bar{F}_V(t, 0)]^2} \\ \bar{R}(t, y) \equiv \overline{\partial_y \bar{F}_V(t, y) \partial_y \bar{F}_V(t, 0)} \end{cases}$$

Correlated disorder (colored-noise):

$$R_\xi(y) = \xi^{-1} R_1(y/\xi)$$

$$\overline{V(t, y)V(t', y')} = D \cdot \delta_{(t-t')} R_\xi(y - y')$$



Free-energy of the $I+I$ DP: 'time'-dependence


$(\xi > 0)$

■ Focus on the two-point correlators:

$$\begin{cases} \bar{C}(t, y) \equiv \overline{[\bar{F}_V(t, y) - \bar{F}_V(t, 0)]^2} \\ \bar{R}(t, y) \equiv \overline{\partial_y \bar{F}_V(t, y) \partial_y \bar{F}_V(t, 0)} \end{cases}$$

■ Correlated disorder (colored-noise):

$$R_\xi(y) = \xi^{-1} R_1(y/\xi)$$



$$\bar{R}(t, y) = \tilde{D}_\infty [\mathcal{R}_\xi(y) - b(t, y, \xi)]$$

Amplitude
 $\tilde{D}_\infty(T, \xi)$

Time dependence:
Encoding the
roughness?

Shape: Microscopic
disorder correlator?

Free-energy of the $1+1$ DP: full evolution

- Stochastic heat equation for the partition function $\mathcal{Z}_V(t, y)$

$$\begin{cases} \partial_t \mathcal{Z}_V(t, y) = \left[\frac{T}{2c} \partial_y^2 - \frac{1}{T} V(t, y) \right] \mathcal{Z}_V(t, y) \\ \mathcal{Z}_V(0, y) = \delta(y) \end{cases}$$

$$\mathcal{Z}_V(t, y) \equiv e^{-F_V(t, y)/T}$$

- KPZ evolution equation for the total free-energy $F_V(t, y)$

$$\begin{cases} \partial_t F_V(t, y) = \frac{T}{2c} \partial_y^2 F_V(t, y) - \frac{1}{2c} [\partial_y F_V(t, y)]^2 + V(t, y) \\ \mathcal{Z}_V(0, y) = e^{-F_V(0, y)/T} = \delta(y) \quad \text{('sharp wedge' initial condition)} \end{cases}$$

$$F_V(t, y) \equiv F_{V \equiv V}(t, y) + \bar{F}_V(t, y)$$

- Tilt KPZ evolution equation for the disorder free-energy $\bar{F}_V(t, y)$

$$\begin{cases} \partial_t \bar{F}_V(t, y) = \frac{T}{2c} \partial_y^2 \bar{F}_V(t, y) - \frac{1}{2c} [\partial_y \bar{F}_V(t, y)]^2 - \frac{y}{t} \partial_y \bar{F}_V(t, y) + V(t, y) \\ \bar{F}_V(0, y) \equiv 0 \quad \text{(flat initial condition)} \end{cases}$$

Free-energy of the $1+1$ DP: linearized evolution

- Tilt KPZ evolution equation for the disorder free-energy $\bar{F}_V(t, y)$

$$\begin{cases} \partial_t \bar{F}_V(t, y) = \frac{T}{2c} \partial_y^2 \bar{F}_V(t, y) - \frac{1}{2c} [\partial_y \bar{F}_V(t, y)]^2 - \frac{y}{t} \partial_y \bar{F}_V(t, y) + V(t, y) \\ \partial_t \bar{F}_V^{\text{lin}}(t, y) = \frac{T}{2c} \partial_y^2 \bar{F}_V^{\text{lin}}(t, y) - \frac{y}{t} \partial_y \bar{F}_V^{\text{lin}}(t, y) + V(t, y) \end{cases}$$

$$F_V^{\text{lin}}(t, y) \equiv F_{V \equiv 0}(t, y) + \bar{F}_V^{\text{lin}}(t, y)$$

- KPZ evolution equation for the total free-energy $F_V(t, y)$

$$\begin{cases} \partial_t F_V(t, y) = \frac{T}{2c} \partial_y^2 F_V(t, y) - \frac{1}{2c} [\partial_y F_V(t, y)]^2 + V(t, y) \\ \partial_t F_V^{\text{lin}}(t, y) = \left[\frac{T}{2c} \partial_y^2 - \frac{y}{t} \partial_y \right] F_V^{\text{lin}}(t, y) + \frac{cy^2}{2t^2} + V(t, y) \end{cases}$$

$$\mathcal{Z}_V^{\text{lin}}(t, y) \equiv e^{-F_V^{\text{lin}}(t, y)/T}$$

- Stochastic heat equation for the partition function $\mathcal{Z}_V(t, y)$

$$\begin{cases} \partial_t \mathcal{Z}_V(t, y) = \left[\frac{T}{2c} \partial_y^2 - \frac{1}{T} V(t, y) \right] \mathcal{Z}_V(t, y) \\ \partial_t \mathcal{Z}_V^{\text{lin}}(t, y) = \left\{ \frac{T}{2c} \partial_y^2 - \frac{1}{T} \left[V(t, y) + \frac{1}{2c} (\partial_y \bar{F}_V^{\text{lin}}(t, y))^2 \right] \right\} \mathcal{Z}_V^{\text{lin}}(t, y) \end{cases}$$


Free-energy of the $I+I$ DP: linearized evolution

$$\partial_t \bar{F}_V(t, y) = \frac{T}{2c} \partial_y^2 \bar{F}_V(t, y) - \frac{1}{2c} [\partial_y \bar{F}_V(t, y)]^2 - \frac{y}{t} \partial_y \bar{F}_V(t, y) + V(t, y)$$


- Fluctuations are exactly Gaussian at all 'times' \Rightarrow fully characterized by: $\bar{R} = \overline{\partial \bar{F} \partial \bar{F}}$

$$\bar{R}^{\text{lin}}(t, y) = \frac{cD}{T} [R_\xi(y) - b^{\text{lin}}(t, y, \xi)]$$

$\bar{C}^{\text{lin}}(t, y)$ $\bar{R}^{\text{lin}}(t, y)$ $\frac{1}{2} \partial_y \bar{C}^{\text{lin}}(t, y)$



$\bar{C}^{\text{lin}}(t, y)$ $\bar{R}^{\text{lin}}(t, y)$ $\frac{1}{2} \partial_y \bar{C}^{\text{lin}}(t, y)$



- Asymptote:




$$\bar{R}^{\text{lin}}(\infty, y) = \frac{cD}{T} R_\xi(y)$$

- Scaling with the diffusive roughness:





$$b^{\text{lin}}(t, y, \xi) = \frac{\tilde{b}(y/\sqrt{B_{\text{th}}(t)}, \xi/\sqrt{B_{\text{th}}(t)})}{\sqrt{B_{\text{th}}(t)}}$$

Numerics: 'time'-evolution of the free-energy correlators $(\xi > 0)$

$$\begin{cases} \bar{C}(t, y) \equiv \overline{[\bar{F}_V(t, y) - \bar{F}_V(t, 0)]^2} \\ \bar{R}(t, y) \equiv \overline{\partial_y \bar{F}_V(t, y) \partial_y \bar{F}_V(t, 0)} \end{cases}$$




Numerics: 'time'-evolution of the free-energy correlators $(\xi > 0)$





- Asymptotic disorder free-energy correlator

$$\bar{R}_{\text{sat}}(y) \approx \bar{R}(\infty, y) = \frac{1}{2} \partial_y^2 \bar{C}(\infty, y)$$

- Shape reminiscent of the microscopic disorder correlator used in our numerical study!



$$R_{\xi}(y) \propto \overline{V(0, y)V(0, 0)}$$



E. Agoritsas, V. Lecomte & T. Giamarchi, *Phys. Rev. B* **82**, 184207 (2010).


E. Agoritsas, V. Lecomte & T. Giamarchi, *Phys. Rev. E* **87**, 042406 & 062405 (2013).

Numerics: temperature dependence of the free-energy

$(\xi > 0)$

■ Amplitude of the correlator / Maximum value

$T \approx 0$
 $\tilde{D}_\infty \sim \frac{cD}{T_c}$
 $\mathcal{R}_\xi \approx ??$



$\xi \approx 0$

$\tilde{D}_\infty \lesssim \frac{cD}{T}$


$\mathcal{R}_\xi \approx \mathcal{R}_\xi$

E. Agoritsas, V. Lecomte & T. Giamarchi, *Phys. Rev. B* **82**, 184207 (2010).

E. Agoritsas, V. Lecomte & T. Giamarchi, *Phys. Rev. E* **87**, 042406 & 062405 (2013).

Numerics: disorder contribution to the roughness


$(\xi > 0)$



Blue → Red:
Low-T → High-T



$$B(t) \equiv \overline{\langle y(t)^2 \rangle} = B_{\text{thermal}}(t) + B_{\text{dis}}(t)$$

$$\begin{cases} B(t) \stackrel{(t \rightarrow \infty)}{\sim} A_{(c,D,T,\xi)} t^{4/3} \\ A_{(c,D,T,\xi)} \sim (\tilde{D}_{\infty}/c^2)^{2/3} \end{cases}$$



Numerics: temperature dependence of the roughness

$(\xi > 0)$




$$B(t) \equiv \overline{\langle y(t)^2 \rangle} = B_{\text{thermal}}(t) + B_{\text{dis}}(t)$$

$$\begin{cases} B(t) \stackrel{(t \rightarrow \infty)}{\sim} A_{(c,D,T,\xi)} t^{4/3} \\ A_{(c,D,T,\xi)} \sim (\tilde{D}_{\infty}/c^2)^{2/3} \end{cases}$$

Numerics: temperature-dependence of the roughness

$(\xi > 0)$



$$B(t) \equiv \overline{\langle y(t)^2 \rangle} = B_{\text{thermal}}(t) + B_{\text{dis}}(t)$$

$$\begin{cases} B(t) \stackrel{(t \rightarrow \infty)}{\sim} A_{(c,D,T,\xi)} t^{4/3} \\ A_{(c,D,T,\xi)} \sim (\tilde{D}_\infty / c^2)^{2/3} \end{cases}$$

Analytical predictions:

- At low temperature: $T \ll T_c$




$$A_{(c,D,0,\xi)} \sim \xi^{-2/9} T^0$$

- At high temperature: $T \gg T_c$

$$A_{(c,D,T,0)} \sim T^{-2/3} \xi^0$$

Summary




- Study of the interplay between finite temperature & finite width/disorder correlation length ξ
- Effective description at fixed lengthscale: fluctuations of DP free-energy at fixed 'time'



- Regimes in the disorder free-energy fluctuations & roughness
- Crossover in temperature controlled by free-energy amplitude $\tilde{D}_{\infty}(T, \xi)$ & characteristic temperature $T_c(\xi) = (\xi c D)^{1/3}$
- Imprint of the microscopic disorder correlator in free-energy correlator $\bar{R}_{\text{sat}}(y)$




Perspectives

- Theoretical predictions to be challenged with experimental interfaces:
 - Ferromagnetic domain walls? (low temperature)
 - Nematic liquid crystals? (high-velocity)
- Connections with the KPZ universality class: role of a correlated disorder/noise?



Perspectives

- Theoretical predictions to be challenged with experimental interfaces:
 - Ferromagnetic domain walls? (low temperature)
 - Nematic liquid crystals? (high-velocity)
- Connections with the KPZ universality class: role of a correlated disorder/noise?



- Better understanding of the short-'time' regime hidden below t_{sat} & non-Gaussian fluctuations generated by the KPZ nonlinearity

$$\frac{1}{2c} \left[\partial_y \bar{F}_V(t, 0) \right]^2$$

- 
- 🌀 A.-L. Barabási & H. E. Stanley, « *Fractal Concepts in Surface Growth* », Cambridge University Press, (1995).
 - 🌀 T. Giamarchi, « *Disordered Elastic Media* » in *Encyclopedia of Complexity and Systems Science*, pp.2019-2038, ed. Springer (2009).
 - 🌀 « *Disordered Systems* » in *Comptes Rendus de Physique* **14**, 637 (2013), editor: T. Giamarchi.

 - 🌀 E. Agoritsas, V. Lecomte & T. Giamarchi, *Phys. Rev. B* **82**, 184207 (2010).
 - 🌀 E. Agoritsas, V. Lecomte & T. Giamarchi, *Physica B* **407**, 1725 (2012).
 - 🌀 E. Agoritsas, S. Bustingorry, V. Lecomte, G. Schehr & T. Giamarchi, *Phys. Rev. E* **86**, 031144 (2012).
 - 🌀 E. Agoritsas, V. Lecomte & T. Giamarchi, *Phys. Rev. E* **87**, 042406 (2013).
 - 🌀 E. Agoritsas, V. Lecomte & T. Giamarchi, *Phys. Rev. E* **87**, 062405 (2013).
 - 🌀 E. Agoritsas, PhD thesis (2013), <http://archive-ouverte.unige.ch/unige:30031>.