

# On a Liouville type theorem for harmonic maps to convex spaces via Markov chains

By

Kazuhiro KUWAE \*and Karl-Theodor STURM\*\*

## Abstract

We give a Liouville type theorem for harmonic maps from the spaces admitting no constant bounded harmonic functions equipped with the harmonicity of functions to convex spaces in terms of conservative Markov chains and barycenters. No differentiable structures for the domain and the target are assumed.

## §1. Introduction

The purpose of this note is to give a Liouville type theorem for harmonic map to metric space having a convex property in terms of discrete time Markov chains. Liouville type theorems for harmonic maps between complete smooth Riemannian manifolds have been done by many authors including geometers and probabilists. Eells-Sampson [13] proved that any (bounded) harmonic map from a compact Riemannian manifold with positive Ricci curvature into a complete manifold with non-positive curvature is a constant map. Schoen-Yau [58] also proved that any harmonic map with finite energy from a complete smooth

---

2000 Mathematics Subject Classification(s): 31C05, 60J05, 60J10, 53C22, 53C43, 58E20, 58J60.

*CAT(0)-space, CAT(1)-space, k-convex space, regular geodesic ball, Banach space, barycenter, Jensen's inequality, Markov chain, Markovian kernel, subharmonic function, harmonic map, invariant  $\sigma$ -field, (strong) Liouville property:*

The first author was partially supported by a Grant-in-Aid for Scientific Research (C) No. 16540201 from Japan Society for the Promotion of Science and the second author was supported by Sonderforschungsbereich 611.

\*Department of Mathematics, Faculty of Education, Kumamoto University, Kumamoto 860-8555, Japan.

\*\*Institute of für Angewandte Mathematik, Bonn Universität, Wegelerstrasse 6, 53115 Bonn, Germany

Riemannian manifold with non-negative Ricci curvature into a complete manifold with non-positive curvature is a constant map. Cheng [3] showed that any harmonic map with sublinear growth from a complete Riemannian manifold with non-negative Ricci curvature into an Hadamard manifold is a constant map. Hildebrandt-Jost-Widman [22] (see also [23],[24]) proved a Liouville type theorem for harmonic maps into regular geodesic (open) balls in a complete  $C^3$  Riemannian manifold from a simple or compact  $C^1$  Riemannian manifold. Choi [6] showed a Liouville type theorem for harmonic maps from a complete smooth Riemannian manifold with non-negative Ricci curvature into regular geodesic (open) balls in a complete smooth Riemannian manifold. Kendall (cf. [40],[25],[41]) showed that any bounded harmonic maps from complete smooth Riemannian manifold having the Liouville property for harmonic functions into regular geodesic (open) ball is a constant map using a stochastic tool. He called his result “ultimate Liouville property”. By Yau’s gradient estimate for harmonic functions (see [66],[5]), any complete smooth Riemannian manifold with non-negative Ricci curvature admits both the elliptic Harnack inequality and the *strong Liouville property*, that is, any positive (or lower bounded) harmonic function is always constant. (The strong Liouville property follows from the elliptic Harnack inequality, cf. 5.4.5 in [56].) So Kendall’s result generalizes [6]. Tam [65] proved that a harmonic map  $u$  from a Riemannian manifold  $M$  quasi-isometric to a complete smooth Riemannian manifold with non-negative Ricci curvature into an Hadamard manifold  $N$  is a constant map if  $u$  satisfies  $d_N(u(x), u(y)) = o(d_M^\gamma(x, y))$ ,  $x, y \in M$ ,  $\gamma \in ]0, 1]$ .

Sung-Tam-Wang [64] proved that if any bounded harmonic function on a complete smooth Riemannian manifold which is asymptotically constant on each non-parabolic end, then the same assertion holds for bounded harmonic maps to regular geodesic balls extending the Kendall’s result. Stafford [59] also gave a probabilistic proof of the result of [3]. Jin [29] proved another type of Liouville type theorem on harmonic maps from Euclidean space to general Riemannian manifolds in terms of the asymptotic behavior of the  $C^2$ -map. Colding-Minicozzi II [8],[9] gave related results to [3] on harmonic functions. Rigoli-Setti [55] also gave a Liouville type theorem for harmonic maps from a certain Riemannian manifold with pole. Cheng-Tam-Wang [4], extending the earlier result [58], proved that any harmonic maps with finite energy into Hadamard manifolds also have the Liouville property provided any bounded harmonic function is always constant, and its stochastic proof is done by Atsuji [1] in the framework of  $L$ -harmonic maps.

The spaces in the above results are assumed to have differentiable struc-

tures in the strict sense. Our main interest is the theory of harmonic maps between singular geometric objects like Alexandrov spaces. In such spaces we can not expect differentiable structures in the classical sense. There are various papers on harmonic maps between singular spaces from geometric or analytic point of view (see [20],[45],[46],[14],[16],[17],[18],[19],[30],[31],[32],[34],[39],[36],[37],[38],[52],[53],[27]). In particular, Gromov-Schoen [20] established the existence of energy minimizing locally Lipschitz maps from Riemannian manifolds into Bruhat-Tits buildings and gave Corlette's version of Margulis's super-rigidity theorem. After [20], Korevaar-Schoen [45] constructed harmonic maps from domains in Riemannian manifolds into Hadamard spaces as a boundary value problem. The same strategy can be applied between Riemannian polyhedra (see [14]). Very recently, Izeki-Nayatani [27] constructed combinatorial harmonic maps from complexes to Hadamard spaces and applied it to prove a fixed point theorem. So the theory of harmonic maps between singular objects is nowadays significant in view of geometric or analytic point.

However, there have not been systematic approach to Liouville type theorem for harmonic maps between singular objects. To begin with, we start to prove a Liouville theorem for harmonic maps between such spaces in the framework of discrete time Markov processes according to the argument by Kendall [40]. More concretely, the source space  $X$  is only assumed to have conservative Markov chains or Markovian kernel  $P(x, dy)$ , which is necessary to define the notion of harmonicity, so-called the  $P$ -harmonicity. The second author has studied  $P$ -harmonic maps taking values into CAT(0)-spaces (see [60],[61]). We shall treat more general target spaces than CAT(0)-spaces including a CAT(1)-space with diameter strictly less than  $\pi/2$  (e.g. Example 2). The target  $Y$  which we consider is assumed to have an analogy of barycenter defined as in [12],[21], which is an extended notion of the usual barycenter in CAT(0)-space discussed in [33],[45],[60],[61],[62],[63],[7]. In this framework we can establish our Liouville type theorem, that is, if the conservative Markov chain has a Liouville property for harmonic functions, then the same property holds for harmonic maps under the condition that the target space is a proper metric space, CAT(0)-space, or separable Banach space (Theorem 3.1).

## §2. Framework

Throughout this note,  $(Y, d)$  denotes a complete separable metric space.

**Definition 2.1** (Admissible Function). A non-negative finite function  $\Phi$  on  $Y \times Y$  is said to be *admissible* if  $\Phi$  vanishes only on the diagonal, there

exists an upper semi continuous function  $\psi : [0, \infty[ \rightarrow [0, \infty[$  such that  $\psi > 0$  on  $]0, \infty[$ ,  $\psi(0) = 0$  and  $\Phi(x, y) \leq \psi(d(x, y))$  for  $x, y \in Y$ , and for each  $y \in Y$ ,  $x \mapsto \Phi(x, y)$  is upper semi continuous and for each  $x \in Y$ ,  $y \mapsto \Phi(x, y)$  is continuous. The triple  $(Y, d, \Phi)$  is called *admissible* if  $\Phi$  is an admissible function on  $Y \times Y$ .

Let  $\mathcal{P}(Y)$  be the family of probability measures on  $(Y, \mathcal{B}(Y))$ . For an admissible function  $\Phi$ , we set

$$\mathcal{P}^\Phi(Y) := \left\{ \mu \in \mathcal{P}(Y) \mid \int_Y \Phi(w, z) \mu(dw) < \infty \text{ for all } z \in Y \right\}.$$

When  $\Phi(x, y) = d^p(x, y)$ ,  $p \geq 1$ , we write  $\mathcal{P}^p(Y)$  instead of  $\mathcal{P}^\Phi(Y)$ .

**Definition 2.2** ( $\Phi$ -Barycenter). Let  $(Y, d, \Phi)$  be an admissible space. For any  $\mu \in \mathcal{P}^\Phi(Y)$ , a point  $b(\mu) \in Y$  is said to be  $\Phi$ -*barycenter* of  $\mu$  if for each  $z \in Y$ ,

$$\Phi(b(\mu), z) \leq \int_Y \Phi(w, z) \mu(dw) (< \infty). \quad (2.1)$$

*Remark.* Given  $\mu \in \mathcal{P}^\Phi(Y)$ , the  $\Phi$ -barycenter of  $\mu$  is not necessarily unique in general. See Example 6 below for the non-uniqueness of  $\Phi$ -barycenter. Denote by  $B^\Phi(\mu)$  the family of  $\Phi$ -barycenters of  $\mu$ .

There are many examples of admissible spaces  $(Y, d, \Phi)$  admitting  $\Phi$ -barycenters in our sense. Next is a list of examples.

**Example 1** (CAT(0)-Space). A complete separable metric space  $(Y, d)$  is called the *CAT(0)-space* (*Hadamard space*, or *global NPC space*) if for any pair of points  $\gamma_0, \gamma_1 \in Y$  and any  $t \in [0, 1]$  there exists a point  $\gamma_t \in Y$  such that for any  $z \in Y$

$$d^2(z, \gamma_t) \leq (1-t)d^2(z, \gamma_0) + td^2(z, \gamma_1) - t(1-t)d^2(\gamma_0, \gamma_1). \quad (2.2)$$

By definition,  $\gamma := (\gamma_t)_{t \in [0, 1]}$  is the minimal geodesic joining  $\gamma_0$  and  $\gamma_1$ . Any CAT(0)-space is simply connected. Hadamard manifolds, Euclidean Bruhat-Tits buildings (e.g. metric tree), spiders, booklets and Hilbert spaces are typical examples of CAT(0)-spaces (cf. [63]). Let  $(Y, d)$  be a CAT(0)-space. Then the distance function  $d : Y \times Y \rightarrow [0, \infty[$  is convex (Corollary 2.5 in [63]) and Jensen's inequality (Theorem 6.3 in [63]) can be applied to the convex function  $Y \ni w \mapsto d(w, z)$  for each  $z \in Y$ . In this case,  $\Phi$  can be taken to be the distance  $d$  and the  $\Phi$ -barycenter is given by the usual barycenter for  $\mu \in \mathcal{P}^1(Y)$  over  $(Y, d)$  discussed in [63]. Then  $(Y, d, \Phi)$  is an admissible space admitting  $\Phi$ -barycenters.

**Example 2** ( $k$ -Convex Space; cf. Ohta [50]). A complete separable metric space  $(Y, d)$  is called the  $k$ -convex space if  $(Y, d)$  is a geodesic space and for any three points  $x, y, z \in Y$ , any geodesic  $\gamma := (\gamma_t)_{t \in [0,1]}$  in  $Y$  with  $\gamma_0 = x$ ,  $\gamma_1 = y$ , and all  $t \in [0, 1]$ ,

$$d^2(z, \gamma_t) \leq (1-t)d^2(z, x) + td^2(z, y) - \frac{k}{2}t(1-t)d^2(x, y). \quad (2.3)$$

By definition, putting  $z = \gamma_t$ , we see  $k \in ]0, 2]$ . The inequality (2.3) yields the (strict) convexity of  $Y \ni x \mapsto d^2(z, x)$  for a fixed  $z \in Y$ . Any closed convex subset of a  $k$ -convex space is again  $k$ -convex. Every CAT(0)-space is a 2-convex space. It is proved by Ohta [50] that any CAT(1)-space (in particular any spherical building ([2]))  $Y$  with  $\text{diam}(Y) \leq \frac{\pi}{2} - \varepsilon$ ,  $\varepsilon \in ]0, \frac{\pi}{2}[$  is a  $\{(\pi - 2\varepsilon) \tan \varepsilon\}$ -convex space, and any Banach space  $L^p$  with  $p \in ]1, 2]$  over a measurable space is a  $2(p-1)$ -convex space (Propositions 3.1 and 3.4 in [50]). He proved that any two points in a  $k$ -convex space can be connected by a unique minimal geodesic (Lemma 2.2 in [50]) and contractible (Lemma 2.3 and Corollary 2.4 in [50]). In this case,  $\Phi$  can be taken to be the square of distance  $d^2$  and the  $\Phi$ -barycenter is given by the pure barycenter  $\bar{b}(\mu)$  for  $\mu \in \mathcal{P}^2(Y)$  over  $(Y, d)$  discussed in Section 4. Then  $(Y, d, \Phi)$  is an admissible space admitting  $\Phi$ -barycenters.

**Example 3** (Regular Geodesic Ball in Riemmanian Manifold). Let  $(Y, d)$  be the regular geodesic (closed) ball with center  $o$  and radius  $r > 0$  in an  $m$ -dimensional complete smooth Riemmanian manifold  $(M, g)$  ( $m \geq 2$ ), where  $d$  is the Riemmanian distance from  $g$ . That is,  $Y := \{x \in M \mid d(o, x) \leq r\}$  does not intersect the cut-locus of the center  $o \in Y$  and the upper bound  $\kappa (= \kappa(r))$  of sectional curvatures in  $Y$  satisfies  $0 \leq \kappa < (\frac{\pi}{2r})^2$  (see [24],[41]). Then, for  $\kappa > 0$ ,  $\Phi$  given by

$$\Phi(w, z) := \frac{1}{\kappa} \cdot \frac{1 - \cos(\sqrt{\kappa}d(w, z))}{\cos(\sqrt{\kappa}d(w, o))}$$

determines a non-negative bounded convex function  $w \mapsto \Phi(w, z)$  for each  $z \in Y$  and  $\Phi(w, z) \leq \frac{d^2(w, z)}{2 \cos \sqrt{\kappa}r}$  for  $w, z \in Y$  (such form of  $\Phi$  is originally indicated by Jäger-Kaul [28]). The barycenter  $b(\mu)$  for  $\mu \in \mathcal{P}(Y)$  defined as a local minimizer of  $Y \ni z \mapsto \int_Y d^2(z, w)\mu(dw)$  ([41]) is called the *Karcher mean* of  $\mu$  and it is uniquely determined in  $Y$  (see Theorem 7.3 in [41]). The following Jensen's inequality holds (see Lemma 7.2 in [41]): Let  $b(\mu)$  be a Karcher mean of a probability measure  $\mu$  on  $Y$  with compact support contained in the interior of  $Y$ . Then for any bounded convex function  $\varphi$ ,

$$\varphi(b(\mu)) \leq \int_Y \varphi(w)\mu(dw). \quad (2.4)$$

When  $\kappa$  is independent of  $r > 0$ , (2.4) remains true for any  $\mu \in \mathcal{P}(Y)$  and bounded convex function  $\varphi$  (we can construct a regular geodesic ball  $U$  containing  $Y$ ). Hence  $\Phi(b(\mu), z) \leq \int_Y \Phi(w, z)\mu(dw)$  for each  $z \in Y$  if  $\kappa$  is independent of  $r > 0$ . Then  $(Y, d, \Phi)$  is an admissible space admitting  $\Phi$ -barycenters under this assumption.

**Example 4** (Banach Space). Let  $(Y, d)$  be a separable Banach space, where  $d$  is given through the norm on  $Y$ . In this case,  $\Phi$  can be taken to be the distance  $d$  and  $\Phi$ -barycenter can be given by the barycenter  $b(\mu)$  for  $\mu \in \mathcal{P}^1(Y)$  defined as the Bochner integral  $\int_Y x\mu(dx)$ . Then  $(Y, d, \Phi)$  is an admissible space admitting  $\Phi$ -barycenters.

Let  $(X, \mathcal{X})$  be a measurable space. Hereafter, we fix a conservative Markov chain  $\mathbf{M} = (\Omega, X_n, \theta_n, \mathcal{F}_n, \mathcal{F}_\infty, \mathbb{P}_x)_{x \in X}$ . Here  $\Omega := X^{\mathbb{N} \cup \{0\}}$  is the family of sequences  $\omega = \{\omega(n)\}_{n \in \mathbb{N} \cup \{0\}}$ ,  $X_n(\omega) := \omega(n)$ ,  $n \in \mathbb{N} \cup \{0\}$ ,  $\theta_n$  is the shift operator defined by  $X_{m+n}(\omega) = X_m(\theta_n \omega)$ ,  $\mathcal{F}_\infty := \sigma\{X_n \mid n \in \mathbb{N} \cup \{0\}\}$ ,  $\mathcal{F}_n := \sigma\{X_k \mid k \leq n\}$ . Denote by  $P(x, dy) := \mathbb{P}_x(X_1 \in dy)$  the transition kernel of  $\mathbf{M}$ , which is a kernel on  $(X, \mathcal{X})$  and set  $Pf(x) := \int_X f(y)P(x, dy) = \mathbb{E}_x[f(X_1)]$  if the integration has a meaning for an  $\mathcal{X}$ -measurable function  $f$  on  $X$ .

**Definition 2.3** ( $P$ -Harmonic Map). Let  $(Y, d, \Phi)$  be an admissible space and consider an  $\mathcal{X}/\mathcal{B}(Y)$ -measurable map  $u : X \rightarrow Y$  satisfying  $u_*P(x, \cdot) \in \mathcal{P}^\Phi(Y)$ . We set

$$Pu(x) := b(u_*P(x, \cdot)).$$

Here  $u_*P(x, \cdot)$  is a Borel measure defined by  $u_*P(x, \cdot)(A) := P(x, u^{-1}(A))$ ,  $A \in \mathcal{B}(Y)$  and we choose a  $\Phi$ -barycenter  $b(u_*P(x, \cdot)) \in B^\Phi(u_*P(x, \cdot))$  of  $u_*P(x, \cdot)$ . An  $\mathcal{X}$ -measurable function  $f : X \rightarrow \mathbb{R}$  is said to be  $P$ -subharmonic if  $f \leq Pf$  on  $X$  and an  $\mathcal{X}/\mathcal{B}(Y)$ -measurable map  $u : X \rightarrow Y$  is called  $P$ -harmonic if  $u = Pu$  on  $X$ .

*Remark.*

- (1) For an  $\mathcal{X}/\mathcal{B}(Y)$ -measurable map  $u : X \rightarrow Y$ ,  $Pu$  depends on the choice of  $(Y, d, \Phi)$  and the choice of  $\Phi$ -barycenters. We do not require the  $\mathcal{X}/\mathcal{B}(Y)$ -measurability of  $Pu : X \rightarrow Y$  (sufficient conditions for this measurability are given in [61] if  $Y$  is a CAT(0)-space). Note that  $P$ -harmonicity of an  $\mathcal{X}/\mathcal{B}(Y)$ -measurable  $u : X \rightarrow Y$  implies the same measurability of  $Pu : X \rightarrow Y$ .
- (2) If the Markov kernel  $P$  is  $m$ -symmetric, i.e.  $\int_X Pfg dm = \int_X fPg dm$  for non-negative  $\mathcal{X}$ -measurable functions  $f, g$ , then we can extend  $P$  on

$L^2(X; m)$ . A map  $u : X \rightarrow Y$  is called (weakly)  $P$ -harmonic if  $Pu = u$   $m$ -a.e. on  $X$ . From [60], under that  $Y$  is CAT(0)-space,  $u$  is (weakly)  $P$ -harmonic if and only if it is a *minimizer of the energy*

$$E[v] := \frac{1}{2} \int_X d_Y(v(x), v(y))^2 P(x, dy) m(dx)$$

among all map  $v : X \rightarrow Y$ . This means that our  $P$ -harmonicity is closely related to classical harmonic maps between Riemannian manifolds. The harmonic maps between singular geometric objects can be defined as a local minimizer of energy functional (see [45],[14]).

**Lemma 2.1.** *Let  $(Y, d, \Phi)$  be an admissible space. If an  $\mathcal{X}/\mathcal{B}(Y)$ -measurable map  $u : X \rightarrow Y$  satisfying  $u_*P(x, \cdot) \in \mathcal{P}^\Phi(Y)$  is  $P$ -harmonic, then  $x \mapsto \Phi(u(x), y_0)$  is  $P$ -subharmonic for each point  $y_0 \in Y$ . Moreover, if the range of  $u$  is bounded, then  $\Phi(u, y_0)$  is also a bounded function.*

*Proof.* By definition of the  $\Phi$ -barycenter of  $u_*P(x, \cdot)$ , we see

$$\Phi(u(x), y_0) = \Phi(Pu(x), y_0) \leq P\Phi(u(\cdot), y_0) < \infty.$$

The last assertion is clear from  $\Phi(u, y_0) \leq \psi(d(u, y_0))$  and the upper-semicontinuity of  $\psi$ .  $\square$

### §3. Liouville Theorem

**Theorem 3.1** (Liouville Property). *Let  $(Y, d, \Phi)$  be an admissible space. We assume one of the following:*

- (1)  $(Y, d)$  is proper, i.e. every bounded closed set is compact.
- (2)  $(Y, d)$  is a CAT(0)-space with  $\Phi = d$  and  $b(\mu)$  is the barycenter for  $\mu \in \mathcal{P}^1(Y)$ .
- (3)  $(Y, d)$  is a separable Banach space with  $\Phi = d$  and  $b(\mu) = \int_Y x \mu(dx)$  for  $\mu \in \mathcal{P}^1(Y)$ .

*Suppose that any bounded  $P$ -harmonic function on  $X$  is always constant. Then the same property holds for any bounded  $P$ -harmonic map.*

*Remark.*

- (1) If  $\mathbf{M}$  is irreducible and recurrent, i.e., every bounded  $P$ -subharmonic function is constant, then one can prove a Liouville type theorem for  $P$ -harmonic map from  $X$  to  $Y$  without assuming the properness of  $Y$  in (1).

- (2) As in Theorem 5 of [40], one can prove a Liouville type theorem for  $Y$ -valued discrete time martingales over a filtered probability space in the framework of admissible space  $(Y, d, \Phi)$  with  $\Phi$ -barycenter (cf. [62],[7] for (discrete or continuous time) martingales taking values in  $\text{CAT}(0)$ -space). Precisely, if  $Y$ -valued martingale  $(Y_n)$  has a non-random limit  $Y_\infty$  as  $n \rightarrow \infty$ , then  $Y_n = Y_\infty = Y_0$  a.s. Its proof is simpler than the proof of Theorem 3.1.
- (3) In our setting, we need the conservativeness of  $\mathbf{M}$  for technical reasons. On account of the conservativeness, we can deduce Theorem 3.1(2) without assuming the properness of  $Y$ . The probabilistic proof of the Kendall's ultimate Liouville theorem between Riemannian manifolds does not work in our case if the target space is a (not necessarily proper)  $\text{CAT}(0)$ -space. So we employ his potentially theoretical method of the proof, which works well without the properness of the target. However, the conservativeness is implicitly assumed in his second proof (see [41]).

**Example 5** (Simple Random Walk on  $\mathbb{Z}$ ). Let  $\mathbf{M} = (\Omega, X_n, \mathbb{P}_x)_{x \in \mathbb{Z}}$  be a simple random walk on  $\mathbb{Z}$ , i.e.,  $P(x, dy) := \mathbb{P}_x(X_1 \in dy) = p_x \delta_{x+1}(dy) + q_x \delta_{x-1}(dy)$  for  $p_x, q_x \in ]0, 1[$  with  $p_x + q_x = 1$ . Any  $P$ -harmonic function  $f$  is given by

$$f(x) = \begin{cases} f(0) + (f(1) - f(0)) \sum_{\ell=1}^x \prod_{k=1}^{\ell-1} \frac{q_x}{p_x} & \text{if } x > 0, \\ f(0) - (f(1) - f(0)) \sum_{\ell=x+1}^0 \prod_{k=\ell}^0 \frac{p_x}{q_x} & \text{if } x < 0, \\ f(0) + (f(1) - f(0))(x-1) & \text{if } p_x = \frac{1}{2} \text{ for all } x \in \mathbb{Z}. \end{cases}$$

If for given  $\varepsilon > 0$ ,  $q_x > (1 + \varepsilon)p_x$  holds except for finite positive integers  $x$ , or  $p_x > (1 + \varepsilon)q_x$  holds except for finite negative integers  $x$ , then any bounded  $P$ -harmonic function  $f$  satisfies  $f(x) \equiv f(0)$ . More strongly, if  $p_x = \frac{1}{2}$  for all  $x \in \mathbb{Z}$ , then any non-negative  $P$ -harmonic function  $f$  satisfies  $f(x) \equiv f(0)$ , namely,  $\mathbf{M}$  possesses the *strong Liouville property*.

A symmetric simple random walk on an infinite weighted graph treated in [47] satisfying elliptic or parabolic Harnack inequality possesses a strong Liouville property (cf. 5.4.5 in [56]). Other examples on (strong) Liouville property of Markov chains can be found in [49],[26],[15] (see also section 5 in [57]).

*Proof of Theorem 3.1.* Suppose that  $u : X \rightarrow Y$  is a bounded  $P$ -harmonic map, that is,  $u = Pu$  and  $\Phi(u, y)$  is a bounded  $P$ -subharmonic function for any point  $y \in Y$  by Lemma 2.1. We may assume that the image of  $u$  is contained in a closed ball  $\bar{B}_R(o)$  for some  $o \in Y$ . We set  $M_y := \sup_{x \in X} \Phi(u(x), y) < \infty$ .



Then  $h_y(x) := M_y - \Phi(u(x), y)$  is a  $P$ -superharmonic function. According to the Riesz decomposition (Theorem 1.4 in [54]), we have the following:

$$h_y = h_{\infty, y} + Gg_y,$$

where  $h_{\infty, y}(x) := \lim_{n \rightarrow \infty} P^n h_y(x)$  is the  $P$ -harmonic part of  $h_y$  and  $Gg_y$  with  $g_y := (I - P)h_y \geq 0$  is the potential part. Here  $G$  is the potential kernel of  $P$  defined by  $G := \sum_{n=0}^{\infty} P^n$ . By assumption,  $h_{\infty, y}$  is a constant  $c_y$ . Consequently, by putting  $d_y := M_y - c_y$ , we have  $d_y - \Phi(u(x), y) = Gg_y(x)$  for all  $x \in X$ , in particular,  $\Phi(u(x), y) \leq d_y$  holds for all  $x \in X$ .

Let  $\{y_j\} \subset Y$  be a countable dense subcollection of  $Y$ . We denote  $g_{y_i}$  by  $g_i$ . For each  $z \in X$ ,  $r > 0$  and  $N \in \mathbb{N}$ , we claim that there exists  $n \in \mathbb{N}$  such that

$$P^n \left( z, \bigcap_{i=1}^N \{x \in X \mid Gg_i(x) \leq r\} \right) > 0. \quad (3.1)$$

If the left hand side of (3.1) vanishes for all  $n \in \mathbb{N}$  with some  $z \in X$ ,  $r > 0$  and  $N \in \mathbb{N}$ , then

$$G(g_1 + g_2 + \cdots + g_N) > r \quad P^n(z, \cdot)\text{-a.s. on } X \text{ for all } n \in \mathbb{N}.$$

Hence, for such  $z \in X$ ,  $P^n G(g_1 + g_2 + \cdots + g_N)(z) > r$  for all  $n \in \mathbb{N}$ , which contradicts  $\lim_{n \rightarrow \infty} P^n Gg_y(z) = 0$  for any  $y \in Y$ . Here we use  $P^n 1 = 1$  on  $X$ .

From (3.1), for each  $z \in X$ ,  $r > 0$  and  $N \in \mathbb{N}$ ,

$$\bigcap_{i=1}^N \{x \in X \mid Gg_i(x) \leq r\} \neq \emptyset,$$

in particular,

$$\bigcap_{i=1}^N \{y \in \bar{B}_R(o) \mid d_{y_i} - \Phi(y, y_i) \leq r\} \neq \emptyset.$$

If  $(Y, d)$  is proper,  $\bar{B}_R(o)$  is compact. When  $(Y, d)$  is a CAT(0)-space with  $\Phi = d$ , we employ the *weak topology* on  $Y$  in terms of the *weak convergence of net* (see [48]). In this case, by Theorem 2.1 in [30],  $\bar{B}_R(o)$  is *sequentially weakly compact*, hence it is *countably weakly compact*, in particular, any decreasing sequence of non-empty closed subsets of  $\bar{B}_R(o)$  has a non-empty intersection (see Definition 1.10 and Remark 1.11 in [11]). Thus, for each  $z \in X$ ,

$$\bigcap_{n=1}^{\infty} \bigcap_{i=1}^{\infty} \{y \in \bar{B}_R(o) \mid d_{y_i} - \Phi(y, y_i) \leq 1/n\} \neq \emptyset.$$

That is, there exists  $y_0 \in \bar{B}_R(o)$  which may depend on  $z$  such that

$$d_{y_i} - \Phi(y_0, y_i) \leq 1/n \text{ for all } i, n \in \mathbb{N},$$

which yields  $d_{y_i} \leq \Phi(y_0, y_i)$  for each  $i \in \mathbb{N}$ . Therefore,  $\Phi(u(x), y_i) \leq \Phi(y_0, y_i)$  for all  $x \in X$  and  $i \in \mathbb{N}$ , hence  $u(x) \equiv y_0$ .

Next we prove the assertion for the case that  $Y$  is a separable Banach space. Recall that  $\Phi(z, w)$  is given by the distance  $d(z, w) = \|z - w\|_Y$  and  $\Phi$ -barycenter is the barycenter defined by Bochner integrals. Let  $u : X \rightarrow Y$  be a bounded  $P$ -harmonic map. Then for each  $\ell \in Y^*$ , we see  $x \mapsto Y^*\langle \ell, u(x) \rangle_Y$  is a  $P$ -harmonic function by use of a property of Bochner integrals. By assumption,  $Y^*\langle \ell, u(x) \rangle_Y \equiv c_\ell$ . Hence for  $x, y \in X$

$$\|u(x) - u(y)\|_Y = \sup_{\ell \in Y^*, \|\ell\|_{Y^*} \leq 1} Y^*\langle \ell, u(x) - u(y) \rangle_Y = 0. \quad (3.2)$$

Therefore,  $u \equiv x_0$  for some  $x_0 \in Y$ . □

#### §4. Appendix: $k$ -convex space

In this section, we clarify that  $k$ -convex space can be an admissible space.

**Lemma 4.1.** *Let  $(Y, d)$  be a  $k$ -convex space. Then every open or closed ball  $B$  in  $Y$  is always convex. In particular, any closed ball  $B$  is again a  $k$ -convex space.*

*Proof.* Noting the convexity of  $x \mapsto d^2(z, x)$ , the proof is easy. □

We can define the notion of *barycenter* of probability measures on a  $k$ -convex space (cf. also Section 5.3 in [51]):

**Definition 4.1** (Barycenter). For  $\mu \in \mathcal{P}^2(Y)$ , we set

$$\text{Var}(\mu) := \inf_{z \in Y} \int_Y d^2(z, y) \mu(dy) (< \infty)$$

and call it the *variance* of  $\mu$ . If  $\mu \in \mathcal{P}^2(Y)$  for  $z \mapsto \int_Y d^2(z, x) \mu(dx)$  admits a unique minimizer, i.e., there exists a unique  $b(\mu) \in Y$  such that

$$\text{Var}(\mu) = \int_Y d^2(b(\mu), y) \mu(dy), \quad (4.1)$$

then  $b(\mu)$  is called the *barycenter* of  $\mu \in \mathcal{P}^2(Y)$ . For  $\mu \in \mathcal{P}^1(Y)$  and  $w \in Y$ , we consider a function  $F_w$  defined by

$$\begin{aligned} F_w(z) &:= \int_Y (d^2(z, x) - d^2(w, x))\mu(dx) \\ &\leq d(z, w) \int_Y (d(z, x) + d(w, x))\mu(dx) < \infty. \end{aligned} \quad (4.2)$$

If  $Y \ni z \mapsto F_w(z)$  admits a unique minimizer  $b(\mu)$ , then it is also called the *barycenter* for  $\mu \in \mathcal{P}^1(Y)$ . The barycenter for  $\mu \in \mathcal{P}^2(Y)$  is automatically the barycenter for  $\mu \in \mathcal{P}^1(Y)$ .

For any closed set  $F$ , we denote by  $C(F)$  the closed convex hull of  $F$ , which is the smallest closed convex set containing  $F$ , that is,  $C(F) := \bigcap \mathcal{C}_F$ ,  $\mathcal{C}_F := \{A \mid A \text{ is closed convex and } F \subset A\}$ .

**Definition 4.2** (Pure Barycenter). Take  $\mu \in \mathcal{P}^1(Y)$  and  $w \in Y$ . If  $C(\text{supp}[\mu]) \ni z \mapsto F_w(z)$  admits a unique minimizer  $\bar{b}(\mu) \in C(\text{supp}[\mu])$ , then it is called the *pure barycenter* of  $\mu \in \mathcal{P}^1(Y)$ .

The following lemma can be similarly proved as in the case of CAT(0)-space.

**Lemma 4.2.** *Let  $(Y, d)$  be a  $k$ -convex space. Then every  $\mu \in \mathcal{P}^1(Y)$  admits both the barycenter  $b(\mu)$  and the pure barycenter  $\bar{b}(\mu)$ .*

*Remark.* Let  $(Y, d)$  be a  $k$ -convex space.

- (1) The pure barycenter  $\bar{b}(\mu)$  of  $\mu \in \mathcal{P}^1(Y)$  is nothing but the barycenter of  $\mu \in \mathcal{P}^1(C(\text{supp}[\mu]))$  over the  $k$ -convex space  $(C(\text{supp}[\mu]), d)$ . If  $b(\mu) \in C(\text{supp}[\mu])$ , in particular, if  $\mu$  has full support, then  $\bar{b}(\mu) = b(\mu)$ .
- (2) Let  $K$  be a closed convex subset of  $(Y, d)$ . The inequality (2.3) yields that there exists a unique minimizer of  $K \ni y \rightarrow d^2(y, x)$  denoted by  $\pi_K x$ , called the *foot-point* or *projected point* to  $K$  from  $x$  and we call the map  $\pi_K : Y \rightarrow K$  the *convex projection* to  $K$ . For  $\mu \in \mathcal{P}^1(Y)$ , we set  $K = C(\text{supp}[\mu])$ . Assume that  $\pi_K$  is contractive, that is,  $d(\pi_K x, \pi_K y) \leq d(x, y)$  for all  $x, y \in Y$ . Then we have  $\bar{b}(\mu) = b(\mu)$ . Indeed, for  $w, z \in Y$  we see

$$\begin{aligned} \int_Y (d^2(\bar{b}(\mu), y) - d^2(w, y))\mu(dy) &\leq \int_K (d^2(\pi_K z, \pi_K y) - d^2(w, y))\mu(dy) \\ &\leq \int_Y (d^2(z, y) - d^2(w, y))\mu(dy). \end{aligned}$$

From Proposition 2.6 in [63], if  $(Y, d)$  is a CAT(0)-space, then  $\pi_K$  is always contractive for any closed convex  $K$ , hence  $\bar{b}(\mu) = b(\mu)$  for  $\mu \in \mathcal{P}^1(Y)$ .

- (3) Let  $(Y, d)$  be a  $k$ -convex space. For  $x_0, x_1 \in Y$  and  $t \in [0, 1]$ ,  $\bar{b}((1-t)\delta_{x_0} + t\delta_{x_1}) = b((1-t)\delta_{x_0} + t\delta_{x_1}) = \gamma_t$ , where  $\gamma_t$  is the point at  $t$  of the minimal geodesic  $\gamma$  joining  $x_0 = \gamma_0$  and  $x_1 = \gamma_1$ . Indeed,  $C(\text{supp}[\mu])$  is the minimal geodesic joining  $x_0$  and  $x_1$  and we see for  $\mu := (1-t)\delta_{x_0} + t\delta_{x_1}$  and any  $z \in Y$ ,

$$\begin{aligned} \int_Y d^2(\gamma_t, y)\mu(dy) &= (1-t)d^2(\gamma_t, \gamma_0) + td^2(\gamma_t, \gamma_1) \\ &= (1-t)t^2d(x_0, x_1) + t(1-t)^2d^2(x_0, x_1) \\ &= t(1-t)d^2(x_0, x_1) \leq (1-t)d^2(z, x_0) + td^2(z, x_1) \\ &= \int_Y d^2(z, y)\mu(dy). \end{aligned}$$

- (4) Let  $(Y, d)$  be a 2-dimensional  $\ell^p$  space  $(\mathbb{R}^2, \|\cdot\|_p)$ ,  $p \in ]1, 2[$ . Then  $(Y, d)$  is a  $2(p-1)$ -convex space by [50]. In this case,  $b(\mu) \neq \int_Y x\mu(dx)$  and  $\bar{b}(\mu) \neq \int_Y x\mu(dx)$  for some  $\mu \in \mathcal{P}^1(Y)$ , where  $\int_Y x\mu(dx)$  is the Bochner integral (see Example 6 below).

**Proposition 4.1** (Jensen's Inequality). *Suppose that  $(Y, d)$  is a  $k$ -convex space. Let  $\varphi$  be a lower semi continuous convex function on  $Y$  and take  $\mu \in \mathcal{P}^1(Y)$ . Then we have*

$$\varphi(\bar{b}(\mu)) \leq \int_Y \varphi d\mu \quad (4.3)$$

*provided the right hand side is well-defined.*

*Proof.* We may assume  $Y = C(\text{supp}[\mu])$  replacing  $Y$  with  $C(\text{supp}[\mu])$ . Under this assumption, pure barycenter can be treated as the barycenter. Then the proof is quite same as in the proof for the case of CAT(0)-spaces (see Theorem 6.2 in [63]).  $\square$

By Proposition 4.1, we can see that every  $k$ -convex space  $(Y, d)$  is an admissible space admitting  $\Phi$ -barycenter by taking  $\Phi = d^2$  with the notion of pure barycenters. Finally, we show that  $\Phi$ -barycenter of a probability measure on  $(Y, \mathcal{B}(Y))$  is not unique over a  $k$ -convex space  $(Y, d)$ .

**Example 6.** Let us consider a 2-dimensional  $\ell^p$  space  $(\mathbb{R}^2, d_{\ell^p})$ ,  $p \in ]1, 2[$  with  $d_{\ell^p}(x, y) := \|x-y\|_p = (|x_1-x_2|^p + |y_1-y_2|^p)^{1/p}$ ,  $x = (x_1, x_2)$ ,  $y = (y_1, y_2)$ . Take three points  $0 = (0, 0)$ ,  $e_1 := (1, 0)$ ,  $e_2 = (0, 1)$  in  $\mathbb{R}^2$  and a probability measure  $\mu := \frac{1}{3}\delta_{e_1} + \frac{1}{3}\delta_0 + \frac{1}{3}\delta_{e_2}$ , and let  $\Delta 0e_1e_2$  be the closed convex set

surrounded by the geodesic triangle among  $0, e_1, e_2$ . Note that any line segment between two points in  $\mathbb{R}^2$  is the minimal geodesic in  $(\mathbb{R}^2, d_{\ell^p})$  joining them. We show  $b(\mu) \neq \int_{\mathbb{R}^2} x\mu(dx)$  and  $\bar{b}(\mu) \neq \int_{\mathbb{R}^2} x\mu(dx)$ . For  $(x_1, x_2) \in \mathbb{R}^2$ , set

$$f_p(x_1, x_2) := (|1 - x_1|^p + |x_2|^p)^{\frac{2}{p}} + (|x_1|^p + |x_2|^p)^{\frac{2}{p}} + (|x_1|^p + |1 - x_2|^p)^{\frac{2}{p}}.$$

Then  $b(\mu) \in \mathbb{R}^2$  (resp.  $\bar{b}(\mu) \in \Delta 0e_1e_2$ ) is the unique minimizer of  $\mathbb{R}^2 \ni (x_1, x_2) \mapsto f_p(x_1, x_2)$  (resp.  $\Delta 0e_1e_2 \ni (x_1, x_2) \mapsto f_p(x_1, x_2)$ ). On the other hand, we see  $\int_{\mathbb{R}^2} x\mu(dx) = (1/3, 1/3)$ . Since

$$f_p\left(\frac{1}{3}, \frac{1}{3}\right) = \frac{2(2^p + 1)^{2/p} + 2^{2/p}}{9} \rightarrow \frac{22}{9} > 2 = f_p(0, 0) \geq f_p(\bar{b}(\mu)) \geq f_p(b(\mu))$$

as  $p \rightarrow 1$ , we have  $b(\mu) \neq (1/3, 1/3)$  and  $\bar{b}(\mu) \neq (1/3, 1/3)$  for some  $p \in ]1, 2[$ . We consider  $\Phi(x, y) := \|x - y\|_p^2$  for  $x, y \in \mathbb{R}^2$ . Then  $(\mathbb{R}^2, d_{\ell^p}, \Phi)$  is an admissible space. But we have two kinds of different  $\Phi$ -barycenters for such  $\mu$  and  $p$ . One is  $\bar{b}(\mu)$ , another is  $\int_{\mathbb{R}^2} x\mu(dx)$ .

## References

- [1] Atsuji, A., Parabolicity, the divergence theorem for  $\delta$ -subharmonic functions and applications to the Liouville theorems for harmonic maps, *Tohoku Math. J. (2)* **57** (2005), no. 3, 353–373.
- [2] Bridson, M. R. and Haefliger, A., *Metric spaces of non-positive curvature*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. **319**, Springer-Verlag, Berlin, 1999.
- [3] Cheng, S. Y., Liouville theorem for harmonic maps. Geometry of the Laplace operator (Proc. Sympos. Pure Math., Univ. Hawaii, Honolulu, Hawaii, 1979), pp. 147–151, Proc. Sympos. Pure Math., XXXVI, Amer. Math. Soc., Providence, R.I., 1980.
- [4] Cheng, S. Y. Tam, L.-F. and Wan, T. Y.-H., Harmonic maps with finite total energy, *Proc. Amer. Math. Soc.*, **124** (1996), 275–284.
- [5] Cheng, S. Y. and Yau, S. T., Differential equations on Riemannian manifolds and their geometric applications, *Comm. Pure Appl. Math.* **28** (1975), no. 3, 333–354.
- [6] Choi, H. I., On the Liouville theorem for harmonic maps, *Proc. Amer. Math. Soc.* **85** (1982), no. 1, 91–94.
- [7] Christiansen, T. and Sturm, K.-Th., Expectations and martingales in metric spaces, preprint, 2005, to appear in *Stochastics*.
- [8] Colding, T. H. and Minicozzi II, W. P., Generalized Liouville properties of manifolds, *Math. Res. Lett.*, **3** (1996), no. 6, 723–729.
- [9] Colding, T. H. and Minicozzi II, W. P., Harmonic functions with polynomial growth, *J. Differential Geom.*, **45** (1997), no. 1, 1–77.
- [10] Corcuera, J. M. and Kendall, W. S., Riemannian barycentres and geodesic convexity, *Math. Proc. Cambridge Philos. Soc.* **127** (1999), no. 2, 253–269.
- [11] G. Dal Maso, *An introduction to  $\Gamma$ -convergence*, Birkhäuser Boston Inc., Boston, MA, 1993.
- [12] Doss, S., Sur la moyenne d'un élément aléatoire dans un espace distancié. (French) *Bull. Sci. Math. (2)* **73**, (1949). 48–72.

- [13] Eells, J. Jr. and Sampson, J. H., Harmonic mappings of Riemannian manifolds, *Amer. J. Math.* **86** 1964 109–160.
- [14] Eells, J. Jr. and Fuglede, B., *Harmonic maps between Riemannian polyhedra*. (English summary) With a preface by M. Gromov. Cambridge Tracts in Mathematics, **142**. Cambridge University Press, Cambridge, 2001.
- [15] Erickson, K. B., The strong Liouville property for a class of random walks, *Monatsh. Math.* **109** (1990), no. 3, 237–246.
- [16] Fuglede, B., Hölder continuity of harmonic maps from Riemannian polyhedra to spaces of upper bounded curvature, *Calc. Var. Partial Differential Equations* **16** (2003), no. 4, 375–403.
- [17] Fuglede, B., Finite energy maps from Riemannian polyhedra to metric spaces, *Ann. Acad. Sci. Fenn. Math.* **28** (2003), no. 2, 433–458.
- [18] Fuglede, B., The Dirichlet problem for harmonic maps from Riemannian polyhedra to spaces of upper bounded curvature, *Trans. Amer. Math. Soc.* **357** (2005), no. 2, 757–792 (electronic).
- [19] Fuglede, B., Dirichlet problems for harmonic maps from regular domains, *Proc. London Math. Soc.* (3) **91** (2005), no. 1, 249–272.
- [20] Gromov M. and Schoen R. M., Harmonic maps into singular spaces and  $p$ -adic super-rigidity for lattices in groups of rank one, *Inst. Hautes Études Sci. Publ. Math.* **76** (1992), 165–246.
- [21] Herer, W., Espérance mathématique au sens de Doss d’une variable aléatoire à valeurs dans un espace métrique. (French) [Mathematical expectation in the sense of Doss of a random variable with values in a metric space] *C. R. Acad. Sci. Paris Sér. I Math.*, **302** (1986), no. 3, 131–134.
- [22] Hildebrandt, S., Jost, J. and Widman, K.-O., Harmonic mappings and minimal submanifolds, *Invent. Math.* **62** (1980/81), no. 2, 269–298.
- [23] Hildebrandt, S., Liouville theorems for harmonic mappings, and an approach to Bernstein theorems. Seminar on Differential Geometry, pp. 107–131, *Ann. of Math. Stud.*, **102**, Princeton Univ. Press, Princeton, N.J., 1982.
- [24] Hildebrandt, S., Harmonic mappings of Riemannian manifolds. Harmonic mappings and minimal immersions (Montecatini, 1984), 1–117, *Lecture Notes in Math.*, **1161**, Springer, Berlin, 1985.
- [25] Huang, H. and Kendall, W. S., Correction note to: “Martingales on manifolds and harmonic maps” [in *Geometry of random motion* (Ithaca, NY, 1987), 121–157, *Contemp. Math.*, **73**, Amer. Math. Soc., Providence, RI, 1988; MR0954635 (90d:58163)] by Kendall, *Stochastics Stochastics Rep.*, **37** (1991), no. 4, 253–257.
- [26] Ichihara, K., A Liouville property for difference operators, *Japan. J. Math. (N.S.)* **24** (1998), no. 1, 123–137.
- [27] Izeki, H. and Nayatani, S., Combinatorial harmonic maps and discrete-group actions on Hadamard spaces, *Geom. Dedicata* **114** (2005), 147–188.
- [28] Jäger, W. and Kaul, H., Uniqueness and stability of harmonic maps and their Jacobi fields, *Manuscripta Math.*, **28** (1979), no. 1-3, 269–291.
- [29] Jin, Z.-R., Liouville theorems for harmonic maps, *Invent. Math.* **108** (1992), no. 1, 1–10.
- [30] Jost, J., Equilibrium maps between metric spaces, *Calc. Var. Partial Differential Equations*, **2** (1994), 173–204.
- [31] Jost, J., Convex functionals and generalized harmonic maps into spaces of non positive curvature, *Comment. Math. Helvetici*, **70** (1995), 659–673.
- [32] Jost, J., Generalized Dirichlet forms and harmonic maps, *Calc. Var. Partial Differential Equations*, **5** (1997), 1–19.
- [33] Jost, J., Nonpositive curvature: geometric and analytic aspects, *Lectures in Mathematics ETH Zürich*. Birkhäuser Verlag, Basel, 1997.
- [34] Jost, J., Generalized harmonic maps between metric spaces, in “Geometric analysis and the calculus of variations,” 143–174, *Internat. Press*, Cambridge, MA, 1996.
- [35] Jost, J., Nonlinear Dirichlet forms, in “New directions in Dirichlet forms,” 1–47, *AMS*

- and Internat. Press, 1998.
- [36] Kasue A., Convergence of Riemannian manifolds and harmonic maps, preprint, 1999.
  - [37] Kasue A., Convergence of Riemannian manifolds, Laplace operators and energy forms, Proceedings of the Fifth Pacific Rim Geometry Conference (Sendai, 2000), 75–97, *Tohoku Math. Publ.*, **20**, Tohoku Univ., Sendai, 2001.
  - [38] Kasue A., Convergence of metric measure spaces and energy forms [MR1962229]. Selected papers on differential equations and analysis, 79–96, *Amer. Math. Soc. Transl. Ser. 2*, **215**, Amer. Math. Soc., Providence, RI, 2005.
  - [39] Kasue A., Convergence of Riemannian manifolds and Laplace operators. II, *Potential Anal.* **24** (2006), no. 2, 137–194.
  - [40] Kendall, W. S., Martingales on manifolds and harmonic maps, Geometry of random motion (Ithaca, N.Y., 1987), 121–157, *Contemp. Math.*, **73**, Amer. Math. Soc., Providence, RI, 1988.
  - [41] Kendall, W. S., Probability, convexity, and harmonic maps with small image I: Uniqueness and fine existence, *Proc. London Math. Soc.*, (3) **61** (1990), no. 2, 371–406.
  - [42] Kendall, W. S., Convex geometry and nonconfluent  $\Gamma$ -martingales. II. Well-posedness and  $\Gamma$ -martingale convergence, *Stochastics Stochastics Rep.*, **38** (1992), no. 3, 135–147.
  - [43] Kendall, W. S., Probability, convexity, and harmonic maps II: Smoothness via probabilistic gradient inequalities, *J. Func. Anal.*, **126** (1994), 226–257.
  - [44] Kendall, W. S., From stochastic parallel transport to harmonic maps. New directions in Dirichlet forms, 49–115, AMS/IP Stud. Adv. Math., **8**, Amer. Math. Soc., Providence, RI, 1998.
  - [45] Korevaar N. J. and Schoen R. M., Sobolev spaces and harmonic maps for metric space targets, *Comm. Anal. Geom.* **1** (1993), 561–659.
  - [46] Korevaar N. J. and Schoen R. M., Global existence theorems for harmonic maps to non-locally compact spaces, *Comm. Anal. Geom.* **5** (1997), 333–387.
  - [47] Kumagai, T., Heat kernel estimates and parabolic Harnack inequalities on graphs and resistance forms, *Publ. RIMS, Kyoto Univ.* **40** (2004), no. 3, 793–818.
  - [48] Kuwae K. and Shioya T., Variational convergence over metric spaces, preprint 2005, to appear in Trans. A.M.S.
  - [49] Lyons, T., Instability of the Liouville property for quasi-isometric Riemannian manifolds and reversible Markov chains, *J. Differential Geom.* **26** (1987), no. 1, 33–66.
  - [50] Ohta, S.-I., Convexities of metric spaces, *Geom. Dedicata* **125**, (2007), no. 1, 225–250.
  - [51] Ohta, S.-I., Extending Lipschitz and Hölder maps between metric spaces, preprint, 2007. Available from <http://www.math.kyoto-u.ac.jp/~sohta/>
  - [52] Picard, J., The manifold-valued Dirichlet problem for symmetric diffusions, *Potential Anal.* **14** (2001), 53–72.
  - [53] Picard, J., Stochastic calculus and martingales on trees, *Annales Institut Henri Poincaré, Prob. Stat.* **41** (2005), 4, 631–683.
  - [54] D. Revuz, *Markov chains*, Second edition. North-Holland Mathematical Library, **11**. North-Holland Publishing Co., Amsterdam, 1984.
  - [55] Rigoli, M. and Setti, A. G., Energy estimates and Liouville theorems for harmonic maps, *Internat. J. Math.* **11** (2000), no. 3, 413–448.
  - [56] Saloff-Coste, L., *Aspects of Sobolev-type inequalities*, London Mathematical Society Lecture Note Series, **289**. Cambridge University Press, Cambridge, 2002.
  - [57] Saloff-Coste, L., Analysis on Riemannian co-compact covers, (English summary) Surveys in differential geometry. Vol. IX, 351–384, Surv. Differ. Geom., IX, Int. Press, Somerville, MA, 2004.
  - [58] Schoen, R. M. and Yau, S.-T., Harmonic maps and the topology of stable hypersurfaces and manifolds with non-negative Ricci curvature, *Comment. Math. Helv.* **51** (1976), no. 3, 333–341.
  - [59] Stafford, S., A probabilistic proof of S.-Y. Cheng’s Liouville theorem. *Ann. Probab.* **18** (1990), no. 4, 1816–1822.
  - [60] Sturm, K.-Th., Nonlinear Markov operators associated with symmetric Markov kernels

- and energy minimizing maps between singular spaces, *Calc. Var. Partial Differential Equations* **12** (2001), no. 4, 317–357.
- [61] Sturm, K.-Th., Nonlinear Markov operators, discrete heat flow, and harmonic maps between singular spaces, *Potential Anal.* **16** (2002), no. 4, 305–340.
  - [62] Sturm, K.-Th., Nonlinear martingale theory for processes with values in metric spaces of nonpositive curvature, *Ann. Probab.* **30** (2002), no. 3, 1195–1222.
  - [63] Sturm, K.-Th., Probability measures on metric spaces of nonpositive curvature. Heat kernels and analysis on manifolds, graphs, and metric spaces (Paris, 2002), 357–390, *Contemp. Math.*, **338**, Amer. Math. Soc., Providence, RI, 2003.
  - [64] Sung, C. J. , Tam, L.-F. and Wang, J., Bounded harmonic maps on a class of manifolds, *Proc. Amer. Math. Soc.* **124** (1996), no. 7, 2241–2248.
  - [65] Tam, L.-F., Liouville properties of harmonic maps, *Math. Res. Lett.*, **2** (1995), no. 6, 719–735.
  - [66] Yau, S.-T., Harmonic functions on complete Riemannian manifolds, *Comm. Pure Appl. Math.* **28** (1975), 201–228.