Problem 1 (Self similarity, 4 points).

Find the parameters α, β such that the differential equations

$$u_t = u_{xx} + (u^2)_x$$

admits a similarity solution in the form $u(x,t) = t^{\alpha} f(\frac{x}{t^{\beta}})$. Find a differential equation for f and show that this ODE can be reduced to first order.

Problem 2 (A nonlinear diffusion equation, 8 points).

Consider the nonlinear diffusion equation

$$u_t - (uu_x)_x = 0 \qquad (x,t) \in \mathbb{R} \times (0,\infty)$$

subject to the initial condition of a unit point source applied at x = 0 at time t = 0, that is, $u(x,0) = \delta(x)$ (where δ is the Dirac delta), with

$$\int_{\mathbb{R}} u(x,t) \, \mathrm{d}x = 1 \,, \qquad \lim_{x \to \pm \infty} u(x,t) = 0 \tag{1}$$

for all t > 0. We look for a self similar solution

$$u(x,t) = t^{\alpha}w(xt^{\beta}).$$

- a) Determine the values of the parameters α, β and the equation solved by w. Hint: impose also the first condition in (1).
- b) Show that the equation for w can be integrated once, to get the ODE

$$3w(y)w'(y) + yw(y) = 0.$$

Obtain a piecewise smooth solution satisfying also (1). Hint: you can use the fact that, because the solution must be symmetric around 0, $u_x(0,t) = 0$ and hence w'(0) = 0.

c) Go back to the original variables and write down a piecewise solution u(x,t). Sketch the function $u(\cdot,t)$ for a few values of t.

Problem 3 (Self-similarity in Burger's equation, 4 points).

Consider Burger's equation

$$u_t + uu_x = 0$$
 $(x, t) \in \mathbb{R} \times (0, \infty)$

with $u(x,0) = \delta(x)$, where δ is the Dirac delta. Recall that by the conservation property

$$\int_{-\infty}^{+\infty} u(x,t) \, \mathrm{d}x = 1 \qquad \text{for all } t \ge 0.$$

Find a (piecewise smooth) self-similar solution. In particular, check that the Rankine-Hugoniot condition is satisfied by your solution. *Hint: the solution should be an N-wave.*

Total: 16 points